



DISCLAIMER

This report has been prepared by the Institute of Geological and Nuclear Sciences Limited (GNS Science) exclusively for and under contract to Taranaki Regional Council. Unless otherwise agreed in writing by GNS Science, GNS Science accepts no responsibility for any use of, or reliance on any contents of this Report by any person other than Taranaki Regional Council and shall not be liable to any person other than Taranaki Regional Council, on any ground, for any loss, damage or expense arising from such use or reliance.

The data presented in this Report are available to GNS Science for other use from August 2011

BIBLIOGRAPHIC REFERENCE

Sherburn, S.; Scott, B.J.; Miller, C.A. 2011. Taranaki Seismicity: July 2010 to June 2011, *GNS Science Consultancy Report* 2011/214. 19p.

CONTENTS

EXECUTIVE SUMMARY	ii
1.0 INTRODUCTION	1
2.0 SEISMOGRAPHS	1
3.0 DATA ANALYSIS	1
3.1 Complete analysis	2
4.0 DATA RELIABILITY	2
5.0 RESULTS	3
6.0 DISCUSSION	4
6.1 Long-term data	4
6.1.1 Larger earthquakes	4
6.1.2 Long-term distribution.....	4
6.1.3 Long-term rate	5
6.2 Volcanic significance	5
6.3 Network effectiveness	5
7.0 CONCLUSIONS	5
8.0 REFERENCES	6

FIGURES

Figure 1.	A map of the seismographs used to locate earthquakes in Taranaki.....	8
Figure 2.	A map of all earthquakes located in Taranaki between July 2010 and June 2011.....	9
Figure 3.	A map of earthquakes less than 50 km deep located in Taranaki between July 2010 and June 2011.	10
Figure 4a.	An east-west cross-section showing earthquakes less than 50 km deep located in Taranaki between July 2010 and June 2011.	11
Figure 4b.	A histogram of the depths of the earthquakes shown in Figure 4a.	11
Figure 5.	A map of earthquakes located in Taranaki between January 1994 and June 2011	12
Figure 6.	A map of earthquakes of M2.7 and larger located in Taranaki between January 1994 and June 2011	13
Figure 7.	A histogram of the number of earthquakes less than 50 km deep that have occurred each month between January 1994 and June 2011 in the area shown in Figure 1	14
Figure 8.	The cumulative number of earthquakes (red line) less than 50 km deep located in the area shown in Figure 1 between January 1994 and June 2011.....	15

ENCLOSURES

CD	- PDF file of GNS Science Consultancy Report 2011/214 Taranaki Seismicity: July 2010 to June 2011	
	- Plus earthquake data	back cover

EXECUTIVE SUMMARY

During the period July 2010 to June 2011, 358 earthquakes were located in the Taranaki region. This accounts for about 2% of the earthquakes located in New Zealand in an average year. Most of the shallow earthquakes occurred in a north-east - south-west trending swath west of Mt Taranaki, and northeast of Stratford. The deeper earthquakes were all located beneath southeast Taranaki. This is similar to the long-term distribution. Nine earthquakes in Taranaki were reported felt.

The largest shallow earthquakes (depth less than 50 km) were four events of magnitude 3.3. One relatively small earthquake was located beneath Mt Taranaki, but we conclude that during the period July 2010 to June 2011 seismic activity on the Taranaki Peninsula was not volcanologically significant.

There were no significant changes at seismographs in Taranaki during the year.

In terms of assessing the effectiveness of the network, technical problems are generally no longer an issue as the new and upgraded sites rarely experience problems. During 2011-12 GNS Science intends to make some minor improvements at two seismographs in Taranaki.

1.0 INTRODUCTION

This report summarises earthquakes in Taranaki for the period 1st July 2010 to 30th June 2011.

The Taranaki Volcano-Seismic Network was commissioned in late-1993, with the first usable data being recorded in January 1994. Since that time data have been recorded almost continuously.

The Taranaki Research Group at Auckland University was responsible for processing data recorded by the network from January 1994 until June 1997. These data were regularly reported to Taranaki Regional Council (TRC), and data from the period January 1994 to March 1995 were summarised in Cavill et al. (1997). In July 1997, GNS Science took over responsibility for data analysis and annual reporting under contract to TRC.

In 2001, the GeoNet project started to improve the existing network for monitoring earthquakes in New Zealand. In 2005, planning began for upgrading seismographs in Taranaki to an equivalent standard to that in other parts of New Zealand. With this upgrade came two important changes. Firstly, the role of the network expanded from solely one of monitoring Mt Taranaki to both monitoring Mt Taranaki and contributing to the New Zealand network as a whole by providing essential earthquake recording capability in the western part of the North Island. Secondly, the concept of a "Taranaki network", with all data sent to, and recorded at, a common point became less clear as multiple data hubs began to be used to send data to GeoNet data centres and sites in Taranaki lost any distinction they may have had from other seismographs in New Zealand. For these reasons we feel that the term "Taranaki Volcano-Seismic Network" no longer has the meaning it used to, and in some cases to refer to a Taranaki-specific network can be both confusing and not convey the information that was intended. For these reasons we are no longer using the term.

2.0 SEISMOGRAPHS

The upgrade to seismographs in Taranaki was completed in 2010 and no changes were made during 2010-11. There are currently nine seismographs in the Taranaki area continuously sending data to GeoNet data centres in Taupo and Wellington for analysis via hubs at the Taranaki Emergency Management Office (TEMO) in New Plymouth, at Kahui Rd west of Mt Taranaki, and at Eltham (Figure 1).

As discussed in the introduction, the distinction between Taranaki seismographs and others in New Zealand has become less in recent years. Several seismographs outside Taranaki often record even relatively small earthquakes that occur in Taranaki and provide data that are often essential to locating these earthquakes. The nearest of these sites are Vera Rd - east of Whangamomona, Hauiti – inland from Awakino, and Wanganui – northwest of Wanganui city.

3.0 DATA ANALYSIS

Seismic data from Taranaki are continuously transmitted to GeoNet data centres in Taupo and Wellington. In near real-time (2-3 minutes), an automatic earthquake detection program is used to search for signals that may be earthquakes. When the signal from an earthquake

is detected, the program attempts to estimate the arrival time of the earthquake waves at each of the seismographs, and those data are saved. Data from all seismographs that may have recorded the earthquake are combined, and using all of these data an attempt is made to automatically locate the earthquake. The automatic location is saved, and if it satisfies certain criteria¹, a message is sent to GeoNet staff pagers, noting the estimated location, time, and magnitude of the earthquake.

When the on-call GeoNet duty scientist receives the pager message they analyse the earthquake data and revise the location. If the earthquake is reported 'felt', or the duty scientist believes it may have been felt, they send the revised location information to the GeoNet web page, an email list (that includes TRC staff), and to GeoNet duty staff pagers.

Each day the GeoNet duty scientists also review all significant earthquakes recorded over the last 24 hours, a process known as 'rapid analysis'. A revised location and magnitude are calculated for these earthquakes. This process is intended to produce a reasonably complete catalogue of the larger New Zealand earthquakes each day, but as all possible earthquakes are not examined, this catalogue is not complete. At this stage any earthquakes reported felt that did not trigger a pager message when they occurred are also sent to the GeoNet web page, an email list, and to GeoNet staff pagers.

The earthquake catalogue is updated as each earthquake is located. This catalogue can be searched via the GeoNet website².

The real-time earthquake analysis system used by GeoNet will change in 2011-12. This is expected to provide more reliable rapid locations and boost GeoNet's capability to respond quickly to significant earthquakes.

3.1 Complete analysis

Since July 2008 complete analysis of seismic data by GeoNet seismic analysts has begun approximately two weeks after data were recorded. This means that the earthquake catalogue is effectively complete and finalised within 2-3 weeks of events occurring. There is currently a period of about 6 months of unprocessed data between September 2007 and June 2008 that is given lower priority and will be cleared more slowly.

4.0 DATA RELIABILITY

The reliability of earthquake locations determined from seismic networks depends on several factors. These include:

- the number of sites at which an earthquake is recorded; small earthquakes are often poorly recorded or recorded on only part of a network and locations for these earthquakes are not as good as those for larger events which are well recorded at many more sites;
- how far an earthquake is from the network; there is little control on the location of earthquakes well outside a network so these are usually poorly located; in the case of

¹ These criteria include position, depth, magnitude and location uncertainty, and are intended to select earthquakes that are felt or are likely to have been felt by a significant number of people.

² <http://magma.geonet.org.nz/resources/quakesearch/>

Taranaki this particularly applies to offshore earthquakes;

- technical problems with a seismic network; these degrade data quality and reduce the number of sites at which an earthquake can be recorded.

If there are insufficient data to locate an earthquake well, it is common practice to fix the depth of that earthquake at some appropriate value, while still calculating the earthquake time and position. In particular, this situation often arises with shallow earthquakes when the nearest seismograph is further from the earthquake than it is deep. Some of the earthquake locations calculated in Taranaki are less reliable than others and have been located with a depth fixed.

During the reporting period there were no known technical problems that caused the loss of data from sites used to locate earthquakes in Taranaki.

5.0 RESULTS

Three hundred and fifty-eight earthquakes were located in the Taranaki region between July 2010 and June 2011 (Figure 2). This amounts to about 2% of earthquakes located in New Zealand in an average year.

Shallow earthquakes (those with a depth less than 50 km) were concentrated in a northeast-southwest trending swath west of Mt Taranaki, with a few north-east of Stratford (Figure 2). The largest shallow earthquakes were four of magnitude 3.3. One small earthquake was located beneath Mt Taranaki. Deep earthquakes (those with a depth greater than 50 km) were mainly located in the Hawera region beneath south-east Taranaki (Figure 2).

Depths were calculated for 80% of the located earthquakes, with the remaining ones fixed at a depth of 5, 12, or 33 km. Earthquakes with a fixed depth were often offshore or were small and less well recorded (Figure 3). Earthquakes for which the depth was not fixed occurred between 5 and about 30 km deep (Figure 4a, b).

Nine earthquakes located in Taranaki (the area shown in Figure 1) were reported felt during the reporting period³. These were:

- 02 August 2010, 21 km depth, offshore Oaonui, magnitude 3.3;
- 07 September 2010, 18 km deep, north-west of Mt Taranaki, magnitude 2.8;
- 02 October 2010, 203 km deep, Hawera area, magnitude 4.9;
- 02 December 2010, 12 km fixed depth, 20 km north-east of Stratford, magnitude 3.3;
- 18 January 2011, 10 km deep, north-west of Mt Taranaki, magnitude 3.3;
- 05 February 2011, 14 km deep, north-west of Mt Taranaki, magnitude 3.0;
- 10 March 2011, 10 km deep, north-west of Mt Taranaki, magnitude 2.8;
- 19 March 2011, 212 km deep, Hawera area, magnitude 4.7;
- 07 April 2011, 135 km deep, Hawera area, magnitude 4.6.

³ Other earthquakes may have been felt, but were not reported. Felt reports can be registered using the GeoNet web site: <http://magma.geonet.org.nz/felt/app?service=external/Felt>.

6.0 DISCUSSION

In this section we compare the seismicity for July 2010 to June 2011 with the earthquake activity in Taranaki since 1994; we discuss any long-term similarities, differences and trends that may be apparent. We also assess the volcanic significance of the recent data and comment on the network effectiveness.

6.1 Long-term data

6.1.1 Larger earthquakes

In earlier reports we have summarised all earthquakes located in Taranaki since January 1994. A seismic network can actually only locate all earthquakes above a certain magnitude, and records only some of the earthquakes of lower magnitude⁴. Summarising all earthquakes may therefore be a little misleading because differences in the distribution may reflect differences in location threshold rather than what earthquakes actually occurred. For an area similar to that shown in Figure 1, but excluding earthquakes more than 20 km offshore, Sherburn and White (2005) showed that for the period 1994 – 2001 the GNS catalogue is complete down to magnitude 2.7.

In summarising data since 1994 we therefore show all located earthquakes in Figure 5 and earthquakes above the M2.7 threshold in Figure 6. Although earthquakes immediately west of Mt Taranaki dominate the catalogue, this is largely a result of the location threshold onshore being significantly lower than that offshore. Plotting only earthquakes of M2.7 and larger (Figure 6), it is clear that the level of activity onshore west of Mt Taranaki is quite similar to that offshore north and south of the Taranaki Peninsula.

A result of showing only earthquakes of M2.7 and larger is that less well-located earthquakes, which tend to be of lower magnitude, are not shown. A plot of earthquakes larger than the location threshold can therefore be one way of showing earthquakes that have more reliable locations. However, by showing only the larger earthquakes we may inadvertently miss some important data. In Figure 5 there are more than 100 earthquakes shown beneath Mt Taranaki, which gives the impression that there is a low, but significant, level of seismicity beneath the volcano, and raises questions about the possible volcanic significance of these events. However, only a few of these earthquakes are large enough to be shown in Figure 6, and many may therefore have locations that are unreliable (particularly those from the 1990s), and possibly mis-located from the more active area immediately west of Mt Taranaki. This does not mean that we should only consider Taranaki earthquakes of M2.7 and larger because a sequence of small earthquakes beneath Mt Taranaki may still have significance as a volcanic precursor.

6.1.2 Long-term distribution

In terms of the distribution of earthquakes, the data for July 2010 to June 2011 (Figure 2) are very similar to those since 1994 (Figures 5 and 6). Both time periods are dominated by a swath of shallow earthquakes (depth less than 50 km) west of Mt Taranaki that trend northeast – southwest, deep earthquakes (depth greater than 50 km) in south-east Taranaki, and a cluster of shallow activity north-east of Stratford.

⁴ This is called the location threshold or magnitude of completeness, and can depend on the area considered, the time interval, and on the analysis procedures used.

The swath of earthquakes west of Mt Taranaki represents part of the Cape Egmont Fault Zone, a zone of seismically active faulting. The deep earthquakes in the Hawera region reflect the bottom of the Pacific plate subducting beneath the North Island. The cluster of earthquakes northeast of Stratford is the western part of a band of activity that continues almost to Mt Ruapehu. All of these are long-term features of the Taranaki seismicity.

6.1.3 Long-term rate

We use two measures to show the long-term, shallow (depth less than 50 km) seismicity in the Taranaki region: the number of located earthquakes each month (Figure 7) and the cumulative number of earthquakes located since 1994 (Figure 8). Both figures show that while there are short-term variations in the rate of activity, the long-term rate of shallow seismicity in the Taranaki region is relatively uniform. Short periods of above average activity occurred in early 2000 (a sequence of earthquakes about 20 km north-east of Stratford), May 2005 to March 2006 (widespread activity, but especially west of Mt Taranaki), and May 2007 (several clusters of activity north and west of Mt Taranaki), and possibly in early 2009 (west and north-west of Mt Taranaki).

6.2 Volcanic significance

Although a few earthquakes are usually located beneath Mt Taranaki most years, they are not thought to be volcanologically significant.

While there are no hard and fast rules for assessing whether an earthquake or group of earthquakes are volcanologically significant, the number of earthquakes, their magnitude and the presence or absence of low-frequency earthquakes (McNutt, 2000) are criteria that are often considered. The earthquakes located beneath Mt Taranaki in 2010-11 are not unusual in any of these respects, and they are therefore not thought to be volcanologically significant.

6.3 Network effectiveness

In the past the effectiveness of the network has been significantly affected by technical problems. However, the now upgraded seismographs are much more reliable and technical problems are now rare. No technical problems had a significant effect on Taranaki data in 2010-11.

During 2011-12 GNS Science intends to try to address some issues with ambient noise at Newhall Rd (NWEZ) and upgrade the sensor at Kahui Hut (KHEZ) from short-period to broadband.

7.0 CONCLUSIONS

Seismic activity on the Taranaki Peninsula continues to be dominated by a swath of earthquakes west of Mt Taranaki, with no significant activity beneath the volcano. No earthquakes have been recorded that might indicate ongoing volcanic processes beneath Mt Taranaki. We therefore conclude that during the period July 2010 to June 2011 seismic activity on the Taranaki Peninsula was not volcanologically significant.

8.0 REFERENCES

- Cavill, A.W., Cassidy, J., Brennan, B.J., 1997. Results from the new seismic monitoring network at Egmont Volcano, New Zealand: tectonic and hazard implications. *New Zealand journal of geology and geophysics* 40: 69-76.
- McNutt, S.R., 2000. Volcano seismicity. In H. Sigurdsson (Editor-in-Chief), *Encyclopaedia of Volcanoes*. Academic Press, pp. 1015-1033.
- Sherburn, S., White, R.S., 2005. Crustal seismicity in Taranaki, New Zealand using accurate hypocentres from a dense network. *Geophysical Journal International* 162: 494-506.

FIGURES




Figure 1. A map of the seismographs used to locate earthquakes in Taranaki. Seismograph sites are indicated by triangles and labelled by their three or four-letter site codes. NEZ is North Egmont, NWEZ is Newall Road, PKE is Pukeiti, DREZ is Durham Road, MHEZ is Mangahewa, NMEZ is Namu Road, KHEZ is Kahui Hut, PREZ is Palmer Road, and LREZ is Lake Rotokare. Hub sites are shown by red circles. Population centres are shaded dark grey and named. The Egmont National Park is shown as a green shaded area. Major roads are shown as grey lines. The inset shows nearby GeoNet seismographs (red dots) that are also used in locating earthquakes in Taranaki.




Figure 2. A map of all earthquakes located in Taranaki between July 2010 and June 2011. Closed circles indicate earthquakes less than 50 km deep and open circles those more than 50 km deep. The size of the symbol is proportional to magnitude of the earthquake, and representative symbols are shown in the top left corner of the diagram. Population centres are shaded dark grey and Egmont National Park is shaded green. Major roads and mapped active faults (thick black lines) are also shown. The lower left inset shows faults that offset the basement rocks but are not considered active faults.




Figure 3. A map of earthquakes less than 50 km deep located in Taranaki between July 2010 and June 2011. Closed circles indicate earthquakes for which a depth was calculated and open circles those for which a depth could not be calculated and was therefore fixed at some appropriate value. The size of the symbol is proportional to the magnitude of the earthquake, and representative symbols are shown in the top left corner of the diagram. Population centres are shaded dark grey and Egmont National Park is shaded green. Major roads and mapped active faults (thick black lines) are also shown. The lower left inset shows faults that offset the basement rocks but are not considered active faults. The dashed box marks the location of the cross-section in Figure 4a.




Figure 4a. An east-west cross-section showing earthquakes less than 50 km deep located in Taranaki between July 2010 and June 2011. The symbols are the same as Figure 3, closed circles for earthquakes for which the depth was calculated and open circles for those for which the depth was fixed. Topography is shown in the upper part of the diagram, with the inverted triangle marking the position of the coast at Cape Egmont. The horizontal: vertical scale is 1:1. The location of the cross-section is shown as a dashed box in Figure 3.

Figure 4b. A histogram of the depths of the earthquakes shown in Figure 4a. Only earthquakes for which the depth was calculated are shown (closed circles in Figure 4a).




Figure 5. A map of earthquakes located in Taranaki between January 1994 and June 2011. Data are from the GNS earthquake database. Closed circles indicate earthquakes less than 50 km deep and open circles those more than 50 km deep. The size of the symbol is proportional to the magnitude of the earthquake, and representative symbols are shown in the top left corner of the diagram. Population centres are shaded dark grey and Egmont National Park is shaded green. Major roads and mapped faults (thick black lines) are also shown. The lower left inset shows faults that offset the basement rocks but are not considered active faults.




Figure 6. A map of earthquakes of M2.7 and larger located in Taranaki between January 1994 and June 2011. Data are from the GNS earthquake database. Closed circles indicate earthquakes less than 50 km deep and open circles those more than 50 km deep. The size of the symbol is proportional to the magnitude of the earthquake, and representative symbols are shown in the top left corner of the diagram. Population centres are shaded dark grey and Egmont National Park is shaded green. Major roads and mapped faults (thick black lines) are also shown. The lower left inset shows faults that offset the basement rocks but are not considered active faults.




Figure 7. A histogram of the number of earthquakes less than 50 km deep that have occurred each month between January 1994 and June 2011 in the area shown in Figure 1. Three separate histograms are shown: all earthquakes (bottom), those of magnitude 3 and above (centre) and those of magnitude 4 and above (top).




Figure 8. The cumulative number of earthquakes (red line) less than 50 km deep located in the area shown in Figure 1 between January 1994 and June 2011. The grey line shows the mean rate.



www.gns.cri.nz

Principal Location

1 Fairway Drive
Avalon
PO Box 30368
Lower Hutt
New Zealand
T +64-4-570 1444
F +64-4-570 4600

Other Locations

Dunedin Research Centre
764 Cumberland Street
Private Bag 1930
Dunedin
New Zealand
T +64-3-477 4050
F +64-3-477 5232

Wairakei Research Centre
114 Karetoto Road
Wairakei
Private Bag 2000, Taupo
New Zealand
T +64-7-374 8211
F +64-7-374 8199

National Isotope Centre
30 Gracefield Road
PO Box 31312
Lower Hutt
New Zealand
T +64-4-570 1444
F +64-4-570 4657